

Signalé par JL. Saladin et transmis par Fr. Héran à Isabelle et le vélo

# Systematic review and meta-analysis evaluating the effects electric bikes have on physiological parameters

Jenna McVicar  | Michelle A. Keske | Reza Daryabeygi-Khotbehsara | Andrew C. Betik | Lewan Parker | Ralph Maddison

Institute for Physical Activity and Nutrition, Deakin University, Geelong, Victoria, Australia

### Correspondence

Jenna McVicar, School of Exercise and Nutrition Sciences, Deakin University, 221 Burwood Highway, Geelong, Vic., 3125, Australia.  
Email: mcvicarj@deakin.edu.au

### Funding information

JM has received full scholarship for completion of PhD from Deakin University

**Background:** There is a universal need to increase the number of adults meeting physical activity (PA) recommendations to help improve health. In recent years, electrically assisted bicycles (e-bikes) have emerged as a promising method for supporting people to initiate and maintain physical activity levels. To the best of our knowledge, there have been no meta-analyses conducted to quantify the difference in physiological responses between e-cycling with electrical assistance, e-cycling without assistance, conventional cycling, and walking.

**Methods:** A systematic review and meta-analysis was conducted following PRISMA guidelines. We identified short-term e-bike studies, which utilized a crossover design comparing physiological outcomes when e-cycling with electrical assistance, e-cycling without electrical assistance, conventional cycling, or walking. Energy expenditure (EE), heart rate (HR), oxygen consumption ( $VO_2$ ), power output (PO), and metabolic equivalents (METs) outcomes were included within the meta-analysis.

**Results:** Fourteen studies met our inclusion criteria ( $N = 239$ ). E-cycling with electrical assistance resulted in a lower energy expenditure (EE) [SMD =  $-0.46$  ( $-0.98, 0.06$ ),  $p = 0.08$ ], heart rate (HR) [MD =  $-11.41$  ( $-17.15, -5.68$ ),  $p < 0.000$ , beats per minute], oxygen uptake ( $VO_2$ ) [SMD =  $-0.57$  ( $-0.96, -0.17$ ),  $p = 0.005$ ], power output (PO) [MD =  $-31.19$  ( $-47.19$  to  $-15.18$ ),  $p = 0.000$ , Watts], and metabolic equivalent (MET) response [MD =  $-0.83$  ( $-1.52, -0.14$ ),  $p = 0.02$ , METs], compared with conventional cycling. E-cycling with moderate electrical assistance resulted in a greater HR response [MD  $10.38$  ( $-1.48, 22.23$ )  $p = 0.09$ , beats per minute], and  $VO_2$  response [SMD  $0.34$  ( $-0.14, 0.82$ )  $p = 0.16$ ] compared with walking.

**Conclusions:** E-cycling was associated with increased physiological responses that can confer health benefits.

Trial Registration: International Prospective Register of Systematic Reviews (PROSPERO) CRD42020203905: [https://www.crd.york.ac.uk/PROSPERO/display\\_record.php?RecordID=203905](https://www.crd.york.ac.uk/PROSPERO/display_record.php?RecordID=203905)

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Scandinavian Journal of Medicine & Science In Sports* published by John Wiley & Sons Ltd.

**KEYWORDS**

electric bike, energy expenditure, heart rate, human physiology, metabolic equivalents, oxygen uptake, Physical activity, power output

## 1 | BACKGROUND

Regular participation in physical activity (PA) is associated with numerous health benefits including the prevention of cardiovascular disease and type 2 diabetes,<sup>1</sup> yet over one quarter of adults worldwide are physically inactive.<sup>2</sup> The World Health Organization (WHO) estimates 27.5% of adults do not meet the recommended level of PA and have subsequently highlighted the need for PA promotion.<sup>2</sup> There is a universal need to increase the number of adults meeting PA recommendations to help improve health on an individual level and to help ease the burden on healthcare systems.<sup>3</sup>

The WHO recommends all adults should undertake 150–300 min of moderate intensity or 75–150 min of vigorous activity per week or some equivalent combination.<sup>2</sup> The most frequently cited reason for not being physically active by individuals is lack of time;<sup>4</sup> therefore, plans and strategies to promote physical activity levels should entail people implementing activity in their everyday life. Walking and cycling for short journeys (termed active transport) has been shown to be effective for achieving recommended levels of PA;<sup>5</sup> however, cycling for transport accounts for 1% of trips in the USA, Canada, and Australia.<sup>6</sup> In recent years, electrically assisted bicycles (e-bikes) have emerged as a promising method for supporting people to increase PA levels via active transport.<sup>7</sup> Compared with conventional cycling, e-bikes help individuals to cycle further and for longer periods of time.<sup>8</sup> E-bikes also offer an ideal opportunity to encourage individuals to include PA into their daily lives by replacing short car trips with cycling.

A 2018 systematic review of 17 studies ( $N = 300$ ) found e-cycling was associated with improved cardiorespiratory fitness and increased PA levels.<sup>9</sup> A 4-week e-cycling intervention found e-bike use increased cardiorespiratory fitness levels and reduced 2-h post oral glucose tolerance test blood glucose levels.<sup>10</sup> Furthermore, another 4-week e-bike intervention found e-cycling increased power output. E-cycling was also associated with feelings of enjoyment among those who were physically inactive.<sup>10</sup> While short-term studies assessed in the systematic review reported on the acute physiological effects of e-cycling compared with conventional cycling and walking, no meta-analysis was conducted to quantify the magnitude of effect on physiological parameters. This study extends upon this previous systematic review by updating the literature and

conducting a meta-analysis to determine the magnitude of physiological response elicited by short-term e-cycling. In doing so, we provide much needed data on the physiological responses associated with e-cycling, which will help inform future programs to promote e-cycling and associated health benefits.

## 2 | MATERIALS AND METHODS

This systematic review and meta-analysis were conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) statement (see Appendix S2 for flow diagram) and was registered with PROSPERO [CRD42020203905]. For this review, e-bikes were defined as electrically assisted bicycles requiring the rider to pedal for assistance to be provided.

Inclusion criteria included acute, single ride comparison studies which assessed e-cycling with assistance compared to e-cycling without assistance, conventional cycling or walking. Acute e-bike studies predominantly measure energy expenditure (EE), heart rate (HR), oxygen uptake ( $VO_2$ ), power output (PO) and report metabolic equivalents (METs). As such, studies which measured these physiological responses were included within this systematic review and meta-analysis. Experimental or observational studies, pre- and post-design, quasi-experimental, randomized, and non-randomized cross-over trials were included. Only papers written in English were included. Assessment of outcomes was laboratory or field based. Studies with adults, adolescents, and children were included.

Exclusion criteria included studies without a comparison or control group, studies that examined environmental effects of e-bike use, and long-term studies ( $\geq 4$ -week intervention period).

Search strategy: Databases searched were PsychINFO, MEDLINE, Embase, ISI Web of Science, CINAHL complete, SPORTDiscuss, Scopus, and PubMed, and were searched from database inception to August 2020. Our full search strategy is available in an additional word document (see Appendix S1).

Researchers (JM and MK) independently screened and reviewed titles and abstracts to identify eligible studies. Researchers (JM, AB, and LP) assessed full-text articles for eligibility. Researchers (JM and RD) independently

performed data extraction and quality assessment for the studies included within the meta-analysis. Any discrepancies were discussed with RM.

**Quality assessment:** The quality assessment tool for quantitative studies developed by the Effective Public Health Practice Project<sup>11</sup> was used to assess the overall rating of included studies according to the following criteria: (1) selection bias, (2) study design, (3) control of confounders, (4) blinding, (5) reliability and validity, and (6) withdrawals and dropout. Studies were rated as strong, moderate, or weak across each criterion, which provided an overall rating of quality. This quality assessment tool is recommended for non-RCT studies.<sup>12</sup>

**Data extraction:** JM and RD extracted data from the papers independently. If papers did not include a full data set, the authors were contacted. Data were extracted using Covidence systematic review software (Veritas Health Innovation, Melbourne, Victoria, Australia), and discrepancies were discussed between researchers. Data extracted included authors and their institution, methodological design, participant population and characteristics, intervention characteristics, and intervention outcomes. Data were synthesized and presented in table format with narrative description.

### 3 | SUMMARY MEASURES

Data extracted from the studies were continuous outcomes; therefore, within the meta-analysis, these data were assessed as either mean difference or standardized mean difference (SMD). EE (kcal) data are presented as SMD. HR data are reported as mean difference as all studies included within the meta-analysis reported HR in beats per minute (BPM).  $VO_2$  (L/min or ml/kg/min) is reported as SMD. PO is reported as mean difference as all studies reported PO in watts (W). METs are reported as mean difference. Nine studies reported METs, four studies were included within the MET meta-analysis and are reported as mean difference. All studies which reported METs are described narratively. For comparisons of SMD, the value was reported as units of standard deviation rather than the units reported for the outcome measured. We considered a SMD of 0.2 as a small effect, 0.5 as a moderate effect, and 0.8 as a large effect.<sup>13</sup>

### 4 | RESULTS

A total of 811 articles were obtained from the initial search and one paper was found through hand searching. After removal of 184 duplicates, 628 studies were title and abstract screened. From the 628 identified, 39 full texts were

reviewed. After full-text review, 14 studies were identified with 12 providing complete data, an additional word document containing the PRISMA flow diagram is available (see Appendix S3). One author did not respond to our request for further information and one author was unable to follow-up on our request; thus, 12 studies were included in the meta-analysis and results for the 14 included studies are presented narratively.

Characteristics of all included studies are presented in Table 1. Of the 14 included studies, six assessed EE,<sup>14-19</sup> 12 assessed HR<sup>14,16,17,18,19,20,21,22,23,24,25,26</sup> five assessed PO,<sup>14,16,17,18,20</sup> eight assessed  $VO_2$ ,<sup>15,16,17,19,20,24,25,27</sup> and nine reported METs.<sup>15,16,17,18,19,20,21,24,27</sup> All studies utilized a crossover design. Eight studies were conducted in Europe (two in France, two in Germany, and one in Norway, Switzerland, The Netherlands, and Belgium), five in the USA, and one in Australia. Sample sizes ranged from 3 to 33. Most studies recruited adults who were healthy, two studies recruited adults who were sedentary<sup>20,24</sup> and one study recruited people with coronary artery disease.<sup>15</sup> Six studies compared e-cycling (with electrical assistance) to conventional cycling,<sup>15,19,20,22,25,27</sup> five studies compared e-cycling with no electrical assistance with e-cycling with electrical assistance.<sup>14,17,18,21,26</sup> Two studies compared e-cycling with electrical assistance to conventional cycling and walking.<sup>16,24</sup> Timing between trials ranged from 2 min to 1-week. Distances rode on the bikes ranged from 46m circuits to 16 km routes. Some studies included varying topography<sup>15,16,20,21,23,26,27</sup> where others maintained a flat route.<sup>14,18,19,22</sup> Throughout many of the studies, participants were advised to cycle at a self-selected pace. Only one study<sup>17</sup> pre-specified target speeds, 16 km/h and 21 km/h; this study also included a ride, which allowed participants to e-cycle at a self-selected pace.

To provide context to the levels of electrical assistance e-bikes provide, we reported the standards from Bosch, an e-bike battery supplier. A low level of electrical assistance provided 40% support, moderate assistance provided 100% support, and a high level of assistance can range from 150% to 250% support. When e-cycling with 100% support, the support provided would match that being produced by the individual cycling, for example 40 W output with 100% support would allow an 80W output, similarly 150% support on a 40 W e-cycle would provide a total of 100 W output.

### 5 | QUALITY ASSESSMENT

The quality assessment tool for quantitative studies was applied (Table 2) and resulted in one strong rating,<sup>15</sup> one moderate rating,<sup>19</sup> and 12 weak ratings.<sup>14,16,17,18,20,21,24,25,26,27</sup> Overall, the methods were

TABLE 1 Study design and participant characteristics of included studies.

First author, year	Study design	Country	Participants	Participant characteristics
Alessio, H. 2021	Randomized crossover	USA	Total 30: Age (years) $26.2 \pm 12.7$ Height (m) $1.8 \pm 0.1$ Body weight (kg) $77.6 \pm 18.4$ BMI ( $\text{kg}/\text{m}^2$ ): $25.1 \pm 4.2$	Healthy adults
Berntsen, S. 2017	Randomized crossover	Norway	Total 8: 23–54 years old.	Healthy adults
Bini, R. 2019	Randomized crossover	Australia	Total 20: Age (years) $40 \pm 15$ Height (cm) $177 \pm 8$ Body weight (kg) $78 \pm 11$	10 postal workers, 10 recreational cyclists
Gojanovic, B. 2011	Crossover	Switzerland	Total 18: Age (years): $35.7 \pm 9.7$ Height (m): $1.70 \pm 0.09$ BMI ( $\text{kg}/\text{m}^2$ ): $24.0 \pm 3.3$ Body weight (kg): $70.1 \pm 13.8$	Sedentary adults
Hall, C. 2019	Convergent mixed methods approach	USA	Total 33: Average age: just under 38 years old.	Experienced mountain bikers
Hansen, D. 2018	Randomized crossover clinical trial	Belgium	Total 15: Patients with CAD. Age (years) $64 \pm 7$	Coronary artery disease patients
Hoj, T. 2018	Crossover	USA	33 participants, average age 22 years old.	Healthy adults
Langford, C. 2017	Semi-crossover	USA	6 females, 11 males. BMI ( $\text{kg}/\text{m}^2$ ): females – 23.1, males – 26.1.	Healthy adults
LaSalle, D. 2017	Crossover	USA	Total 12: Females (mean $\pm$ SE): Age (years) $22 \pm 1$ Height (cm) $171 \pm 2$ Weight (kg) $71.2 \pm 5$ Body fat (%) $23.4 \pm 3.3$ Males (mean $\pm$ SE): Age (years) $25 \pm 1$ Height (cm) $177 \pm 2$ Weight (kg) $87.9 \pm 6$ Body fat (%) $16.8 \pm 1.9$	Healthy active adults
Louis, J. 2012	Randomized crossover	France	Total 20: Two participant groups: Trained v untrained. Trained: Age (years) $38.7 \pm 14.8$ , Height (m) $1.77 \pm 0.06$ , Body weight (kg) $69.2 \pm 5.8$ BMI ( $\text{kg}/\text{m}^2$ ): $22.0 \pm 1.1$ Untrained: Age (years) $28.9 \pm 6.3$ , Height (m) $1.72 \pm 0.07$ , Body weight (kg) $66.1 \pm 14.8$ , BMI ( $\text{kg}/\text{m}^2$ ): $22.2 \pm 3.7$	10 trained adults & 10 untrained adults
Meyer, D. 2014	Crossover	Germany	Total 3 males: Age (years) 25–27, Weight (kg) 71–79, Height (cm) 176–183	Recreational cyclists
Simons, M. 2009	Crossover	The Netherlands	12 Total: Age (years): $52.2 \pm 8.7$ Height (cm) $173.3 \pm 7.6$ BMI ( $\text{kg}/\text{m}^2$ ): $24.5 \pm 2.6$ Body weight (kg): $73.6 \pm 9.7$	Habitually active adults – 7 met PA guidelines
Sperlich, B. 2012	Randomized crossover	Germany	8 females, Age (years) $38 \pm 15$ Body mass (Kg) $71.3 \pm 12.9$ BMI ( $\text{kg}/\text{m}^2$ ): $25.3 \pm 2.1$	Sedentary adults
Theurel, J. 2012	Crossover	France	10 total: 5 females: Age (years) $30 \pm 12$ Height (cm) $163 \pm 2$ Weight (kg) $58 \pm 4$ 5 Males: Age (years) $35 \pm 14$ Height (cm) $177 \pm 9$ Weight (kg) $73 \pm 9$	Healthy adults – moderate PA level

Physiological parameter	Intervention characteristics (What was compared?)	Ride characteristics
Energy expenditure (kcal/hr), % HR max, % VO <sub>2</sub> max	E-bike to conventional bike	E-bike low assist v e-bike moderate assist v conventional bike
VO <sub>2</sub> (%)	E-bike to conventional bike	E-bike assistance self-selected v conventional bike
Energy expenditure (kcal), heart rate (bpm), and power output (W)	E-bike with various electrical assistance	E-bike no electrical assistance v e-bike with electrical assistance
Heart rate, VO <sub>2</sub> (l/min)	E-bike was compared with conventional bike and walking	E-bike high assistance v e-bike standard assistance v conventional bike v walking
Heart rate (bpm)	E-mountain bike compared with conventional mountain bike	E-bike assistance not advised v conventional mountain bike
Energy expenditure (kcal), Mean VO <sub>2</sub> (ml/min)	e-bike v conventional bike	E-bike low assistance v e-bike high assistance v conventional bike
Heart rate average (bpm), heart rate max (bpm)	e-bike v conventional bike	E-bike assistance level not available v conventional bike
Energy expenditure (kcal), heart rate (bpm), power output (W), VO <sub>2</sub>	E-bike was compared to conventional bike and walking	E-bike high electrical assistance v conventional bike v walking
Heart rate max (%), VO <sub>2</sub> max (%)	E-bike with various levels of electrical assistance	E-bike with pedal assist mode v e-bike without electrical assistance
Energy expenditure (kcal), heart rate (bpm), power output (W), VO <sub>2</sub> (ml/kg/min)	E-bike with various assistance levels	E-bike unassisted v e-bike light electrical assistance v e-bike high electrical assistance
Lactate (mmol/L), heart rate (bpm), Borg scale.	E-bike with and without electrical assistance	E-bike with electrical assistance v e-bike without electrical assistance
Energy expenditure (kcal), Heart rate (bpm), Power output (W)	E-bike with no support v e-bike with varying electrical assistance	E-bike No Electrical Assistance v E-bike Eco Electrical Assistance v E-bike Power Support
Heart rate (bpm), VO <sub>2</sub> (ml.kg.min), Mean power output (W)	E-bike v conventional bike	E-bike with assistance v conventional bike
Heart rate (bpm) VO <sub>2</sub> (ml.kg.min)	E-bike v conventional bike	E-bike assisted cycling v conventional bike

TABLE 2 Quality assessment tool for quantitative studies.

Study	Selection bias	Design	Confounders	Blinding	Methods	Drop-outs	Global rating
Alessio, 2021	Strong	Moderate	Weak	Weak	Strong	Strong	Moderate
Berntsen, 2017	Weak	Moderate	Weak	Weak	Strong	Strong	Weak
Bini, 2019	Weak	Moderate	Weak	Weak	Strong	Strong	Weak
Gojanovic, 2011	Weak	Weak	Weak	Weak	Strong	Strong	Weak
Hall, 2019	Moderate	Weak	Weak	Weak	Strong	Weak	Weak
Hansen, 2018	Strong	Strong	Strong	Weak	Strong	Strong	Strong
Hoj, 2018	Moderate	Weak	Weak	Weak	Strong	Strong	Weak
LaSalle, 2017	Weak	Moderate	Weak	Weak	Strong	Strong	Weak
Langford, 2017	Weak	Moderate	Weak	Weak	Moderate	Strong	Weak
Louis, 2012	Weak	Moderate	Strong	Weak	Strong	Weak	Weak
Meyer, 2014	Weak	Weak	Moderate	Weak	Strong	Strong	Weak
Simons, 2009	Weak	Weak	Weak	Weak	Strong	Strong	Weak
Sperlich, 2012	Weak	Strong	Weak	Weak	Strong	Weak	Weak
Theurel, 2012	Weak	Weak	Weak	Weak	Strong	Moderate	Weak

Note: Strong = no weak rating; moderate = one weak rating; weak = two or more weak ratings.

rated as strong. Methods were assessed on validation and reliability of the data collection tools used. All included studies were rated as weak for blinding. Blinding was not included in the global rating due to the nature of the methodologies (i.e., the inability to reliably blind participants to an e-bike). Selection bias and confounders were generally rated as weak. Selection bias was rated on the likelihood participants were representative of the target population. Confounders were rated on the differences between study groups and indication of control for these differences. Studies were classified as strong if they received no weak rating, classified as moderate if they received one weak rating and classified as weak if they received two or more weak ratings. One study<sup>15</sup> was powered to detect a significant difference in calorie expenditure between cycling conditions.

## 6 | OUTCOME MEASURES

### 6.1 | Narrative review – METs

Nine studies<sup>15,16,17,18,19,20,21,24,27</sup> assessed METs with e-cycling with electrical assistance compared with conventional cycling or e-cycling without electrical assistance. Reporting of METs within the studies was inconsistent, many authors reported mean or range of MET values, many studies did not report mean and SD. Four studies were included within the meta-analysis,<sup>15,19,20,24</sup> the included studies compared METs between e-cycling with moderate assistance and conventional cycling. Two

studies used walking as a comparator.<sup>16,24</sup> METs ranged from 3 to 10.9. Overall, e-cycling with electrical assistance was associated with lower MET values compared with its control comparator; however, some e-cycling was associated with moderate (3–6 METs)<sup>16–20</sup> to vigorous (MET range >6 METs) intensity activity.<sup>15,21,24,27</sup> Louis et al<sup>17</sup> only provided ranges for MET data and as such were excluded from Table 3. Louis et al<sup>17</sup> reported all participants cycled at an intensity of at least 6 METs with no electrical assistance. With electrical assistance, Louis et al<sup>17</sup> reported untrained participants e-cycled at intensity of >6 METs; a similar intensity was observed only in trained participants who cycled at 21km/h with moderate electrical assistance. They reported the highest level of electrical assistance was associated with 3 to 6 METs.<sup>17</sup> Langford et al<sup>16</sup> provided an average of the MET values from the three segments included within their studies cycle route, a breakdown of the MET values reported are specified in Table 3.

### 6.2 | Meta-analysis–EE, HR, VO<sub>2</sub>, PO, and METs

Random effect meta-analysis was conducted, heterogeneity was assessed by Chi-square test and reported as  $I^2$ . E-cycling with moderate assistance was compared with conventional cycling for all outcomes. E-cycling with moderate assistance was compared with e-cycling without assistance for EE, HR, VO<sub>2</sub>, and PO. E-cycling with high assistance was compared with conventional cycling for HR and VO<sub>2</sub> and with e-cycling without assistance

TABLE 3 Metabolic Equivalent (METs) means from included studies.

Study	Conventional cycling METs (mean)	E-bike no electrical assistance METs (mean)	E-bike moderate electrical assistance METs (mean)	E-bike high electrical assistance METs (mean)	Walking METs (mean)
Alessio, 2021	6.7		5.8	4.8	
Berntsen, 2017	10.9 <sup>a</sup>		8.5 <sup>a</sup>		
Gojanovic, 2011	8.2		7.3	6.1	6.5
Hansen, 2018	6.4		6.6	6.0	
La Salle, 2017		8.5	8.3		
Langford, 2017 (downhill)	3.9		3.7		3.8
Langford, 2017 (flat)	5.2		4.5		4.1
Langford, 2017 (uphill)	7.6		6.6		5.3
Langford, 2017 (average)	5.8		5.1		4.8
Sperlich, 2012	7.1		5.2		
Simons, 2009		6.1	5.7	5.2	

Note: Average MET values of included studies only mean reported to allow for consistency between studies; SD not always reported by authors.

<sup>a</sup>Authors reported median values.

for EE, HR, and PO. A random effects meta-analysis was conducted between e-cycling with moderate assistance and walking for two outcomes, HR and VO<sub>2</sub>. E-cycling with electrical assistance resulted in an increase in physiological responses assessed; however, changes were lower when compared with conventional cycling or e-cycling with no electrical assistance. HR and VO<sub>2</sub> were higher when e-cycling compared with a walking comparison.

Figures 1–5 represent forest plots for EE, HR, VO<sub>2</sub>, PO & MET data compared with conventional cycling. Appendix S4 contains all forest plots.

Tables 4–8 represent meta-analysis information for each outcome with comparator.

Overall, EE (kcal) increased when e-cycling with electrical assistance; however, values were lower than any comparator; there was a small decrease in EE for e-cycling with moderate electrical assistance compared with conventional cycling (SMD  $-0.46$ , 95% CI:  $-0.98, 0.06$ ,  $p = 0.08$ ), however, this finding was not significant. Although an increase in EE from baseline was observed during all cycling conditions, the largest difference between conditions was observed between e-cycling with high electrical assistance and e-cycling without electrical assistance SMD  $2.66$  (95% CI:  $-4.05, -1.28$ ,  $p = 0.0002$ ).

E-cycling with electrical assistance was associated with an increase in HR response; the response was lower when compared with conventional cycling or e-cycling without electrical assistance. E-cycling with moderate electrical assistance resulted in a difference of  $-11.41$  (95% CI:  $-17.15, -5.68$ ) BPM compared with conventional cycling. Compared with walking, e-cycling with electrical assistance was associated with a higher HR (MD  $10.38$ , 95% CI:  $-1.48, 22.23$ ); however, was not significant  $p = 0.09$ .

VO<sub>2</sub> increased from baseline when e-cycling with electrical assistance, e-cycling without assistance, conventional cycling, and walking. VO<sub>2</sub> was higher when e-cycling with electrical assistance compared to walking; however, this finding was not significant (SMD  $0.34$ , 95% CI:  $-0.14, 0.82$ ,  $p = 0.16$ ). VO<sub>2</sub> when e-cycling with a moderate assistance was lower compared with conventional cycling, SMD  $-0.57$  (95% CI:  $-0.96, -0.17$ ,  $p = 0.005$ ).

PO during e-cycling with moderate electrical assistance was lower, mean difference  $-31.19$  W (95% CI:  $-47.19, -15.18$  W,  $p = 0.0001$ ) compared with conventional cycling. E-cycling with high electrical assistance was associated with the largest difference in PO, mean difference  $-53.71$  W (95% CI:  $-64.09, -43.34$  W,  $p < 0.00$ ).

A small difference is observed between e-cycling with moderate assistance compared with conventional cycling,  $-0.83$  METs (95% CI:  $-1.52, -0.14$ ,  $p = 0.02$ ).

## 7 | DISCUSSION

In this systematic review and meta-analysis, we aimed to quantify the physiological response from e-cycling with electrical assistance when compared with e-cycling without electrical assistance, conventional cycling, or walking. Overall, e-cycling was associated with an increase in physiological responses, equivalent to moderate intensity physical activity. Across a host of physiological parameters (EE, HR, VO<sub>2</sub>, PO, and METs), physiological responses were lower than observed with conventional cycling, or e-cycling without electrical assistance, but generally greater than that observed when walking.




FIGURE 1 Forest plot showing the standardized mean difference in energy expenditure response when using an e-bike with moderate electrical assistance v conventional bike




FIGURE 2 Forest plot showing mean difference in heart rate data for e-bike with moderate electrical assistance v conventional bike




FIGURE 3 Forest plot showing oxygen uptake data for e-bike with moderate electrical assistance v conventional bike




FIGURE 4 Forest plot showing mean difference in power output data for e-bike with moderate electrical assistance v conventional bike

Conventional cycling, e-cycling without assistance, and walking were used as comparators to allow for a realistic comparison of modes of active transport as conventional cycling and walking are well-established modes of active travel. E-cycling without assistance and e-cycling with

high assistance were included as comparators as e-bikes without assistance are heavier than a conventional bike and people may not continuously e-cycle with assistance on. Furthermore, people may choose to e-cycle with a high assistance level.






FIGURE 5 Forest plot showing mean difference in metabolic equivalent data for e-bike with moderate electrical assistance v conventional bike

TABLE 4 Main effects of EE (kcal) response to e-cycling with comparison.

E-bike mode	Comparator	No of studies	SMD (95% CI)	I <sup>2</sup> (%)	p value
Moderate electrical assistance	Conventional bike	2	-0.46 (-0.98, 0.06)	0	0.08
Moderate electrical assistance	E-bike no electrical assistance	3	-1.08 (-1.49, -0.66)	0	<0.00001
High electrical assistance	E-bike no electrical assistance	2	-2.66 (-4.05, -1.28)	72	0.0002

TABLE 5 Main effects of HR response to e-cycling with comparison

E-bike mode	Comparator	No of studies	Mean difference (BPM) (95% CI)	I <sup>2</sup> (%)	p value overall
Moderate electrical assistance	Conventional bike	7	-11.41 (-17.15, -5.68)	42	<0.0001
Moderate electrical assistance	E-bike no electrical assistance	2	-3.41 (-10.98, 4.16)	0	0.38
High electrical assistance	Conventional bike	2	-19.50 (-27.32, -11.68)	0	<0.00001
High electrical assistance	E-bike no electrical assistance	2	-15.77 (-23.25, -8.30)	0	<0.0001
Moderate electrical assistance	Walking	2	10.38 (-1.48, 22.23)	51	0.09

TABLE 6 Main effects of VO<sub>2</sub> response to e-cycling with comparison

E-bike mode	Comparator	No of studies	SMD (95% CI)	I <sup>2</sup> (%)	p value
Moderate electrical assistance	Conventional bike	7	-0.57 (-0.96, -0.17)	41	0.005
Moderate electrical assistance	E-bike no electrical assistance	1	-0.89 (-1.86, 0.99)	53	0.08
High electrical assistance	Conventional bike	2	-1.10 (-1.56, -0.65)	0	<0.00001
Moderate electrical assistance	Walking	2	0.34 (-0.14, 0.82)	0	0.16

## 8 | STRENGTHS AND LIMITATIONS

To the best of our knowledge, this is the first meta-analysis to quantify and compare the short-term physiological effects of e-cycling with more conventional forms of active transport (cycling and walking). We undertook a comprehensive search of short-term e-bike studies that assessed physiological responses. This study provides new data on the physiological responses associated with e-cycling and its potential as a public health initiative for promoting recommended levels of physical activity. A limitation was the lack of data for certain outcome measures. Some results were based on two studies (HR and VO<sub>2</sub> for

walking comparisons), which is required by Cochrane as the minimum for a meta-analysis; however, more studies with similar outcomes and methodologies would substantiate these findings. Moreover, topography varied between studies which could have an impact on outcomes. Our search criteria excluded non-English manuscripts meaning studies could have been missed. Reported I<sup>2</sup> values infer substantial heterogeneity between various included studies, to combat this, we ran random effects models which could have produced wider CI.

Many of the studies within this systematic review and meta-analysis included healthy adult participants;<sup>16,19,22,25,27</sup> however, it was not clear how active these adults were. Two studies recruited sedentary adults,<sup>20,24</sup>

TABLE 7 Main effects of PO response to e-cycling

E-bike mode	Comparator	No of studies	Mean difference (W) (95% CI)	I <sup>2</sup> (%)	p value
Moderate electrical assistance	Conventional bike	2	-31.19 (-47.19, -15.18)	0	0.0001
Moderate electrical assistance	E-bike no electrical assistance	3	-19.63 (-22.42, -16.85)	0	<0.00001
High electrical assistance	E-bike no assistance	2	-53.71 (-64.09, -43.34)	75	<0.00001

TABLE 8 Main effects of MET response to e-cycling

E-bike mode	Comparator	No of studies	Mean difference (MET) (95% CI)	I <sup>2</sup> (%)	p value
Moderate electrical assistance	Conventional bike	4	-0.83 (-1.52, -0.14)	31	0.02

one study recruited both trained and untrained participants,<sup>17</sup> and one study investigated a clinical population cohort, people with coronary artery disease.<sup>15</sup> The heterogeneity between the participant groups may have an impact on the outcomes assessed. The quality assessment ratings of the included studies confirm the need for high quality research, future studies should aim to address these limitations.

## 9 | E-CYCLING

E-cycling encourages people to travel further and for longer periods of time,<sup>8</sup> our findings indicate the use of e-bikes would allow people to achieve recommended levels of PA as e-cycling can elicit moderate intensity activity levels. Our results show there is a small difference in METs between e-cycling with moderate assistance and conventional cycling. While offering the benefits of physical activity, e-bikes are often perceived as easier to ride and reduce concerns about cycling distance and hills.<sup>28</sup> E-bikes give riders greater control over their levels of exertion,<sup>29,30</sup> increase feelings of exercise self-efficacy,<sup>8</sup> and extend the active transport radius (from about 5km on a bicycle to 15km or more).<sup>31</sup> Moreover, a recent study<sup>32</sup> interviewed both e-bike users and non e-bike users and highlighted the future potential for e-bikes to be used as an alternative to public transport, especially in a post-COVID-19 pandemic world. However, the interviews indicated that e-cycling was sometimes perceived as physical inactivity;<sup>32</sup> therefore, education is required to advise this is untrue, and health benefits can be obtained from e-cycling.

Although there is evidence to suggest e-cycling is beneficial for health,<sup>9</sup> cost is a considerable barrier for purchase of e-bikes.<sup>32</sup> To promote e-bike use, effort should be made

to make e-bikes accessible to all individuals similar to “The eBike To Work Scheme”<sup>33</sup> or e-bike hire schemes.<sup>34</sup>

## 10 | COMPARISONS WITH OTHER WORK

Our findings complement previous reviews that support the beneficial effects of e-cycling on physical activity levels and health.<sup>9</sup> Bourne et al.<sup>9</sup> reported HR was lower when e-cycling compared with conventional cycling, which is in agreement with our findings. VO<sub>2</sub> was included within the systematic review and was reported to be lower when compared with conventional cycling or e-cycling without support. Existing literature supports that numerous health benefits are obtained from walking.<sup>35</sup> Results from walking interventions have shown increases in aerobic capacity, reductions in systolic and diastolic blood pressure, reductions in waist circumference, reductions in body fat and body mass index.<sup>35</sup> Results from our meta-analysis demonstrate an increased physiological response from e-cycling compared with walking. Results from the meta-analysis established e-cycling with standard electrical support produced higher HR and VO<sub>2</sub> responses compared with walking. As a result, we can infer a similar, if not increased health benefit may be ascertained from e-cycling as an ongoing PA intervention.

## 11 | FUTURE RESEARCH/IMPLICATIONS

There is a clear gap in the literature, e-cycling should be further explored within clinical populations such as those with type 2 diabetes, pre-diabetes, metabolic syndrome, and people who are sedentary and overweight or obese.

Individuals with these metabolic conditions would benefit from increased levels of PA, associated physiological responses and health benefits,<sup>19</sup> particularly as many people with these metabolic conditions face difficulties maintaining PA levels.<sup>36</sup> Evidence suggests those who are inactive will benefit the most by increasing their PA levels; the health benefits are considerable for those increasing PA levels from sedentary to low levels of activity.<sup>37</sup> E-bikes offer an ideal solution and entry-point for such populations, as they elicit the physiological responses and benefits associated with physical activity without many of the barriers associated with conventional cycling, for example, sweating and needing changing facilities,<sup>38</sup> difficulty cycling uphill, and fear of cycle distance.<sup>39</sup> Further research should aim to understand the best ways in which researchers and behavioral scientists can help support individuals to e-cycle and implement e-bikes into their day-to-day lives.

Future research is necessary to compare the health benefits of e-cycling with motorized transport. By assessing the difference in cardiometabolic risk factors between those who e-cycle regularly compared with people who use motorized transport regularly could provide a clearer understanding of the benefits of e-cycling. Furthermore, as societies move towards greener choices,<sup>40</sup> e-cycling is an option that could be utilized due to the health and well-being benefits previously outlined.

## 12 | PERSPECTIVES

Physical inactivity (performing little or no physical activity) is an increasing problem globally. Many adults worldwide do not meet recommended guidelines for physical activity levels and therefore public health strategies are urgently needed to help people engage in regular physical activity. Research has shown that electric bikes, which provide electrical assistance while cycling may offer an ideal approach for increasing physical activity levels, particularly for those who have difficulty exercising. However, little is known about the short-term physiological effects of electric assisted cycling (e-cycling) and how it compares with conventional cycling. Results from this meta-analysis showed that e-cycling was comparable with moderate intensity physical activity, which offer health benefits. Randomized controlled trials are warranted to test this. In summary, e-cycling offers a viable approach to support people to be more physically active. Healthcare professionals might consider encouraging e-cycling when providing options to support people to be physically active. Moreover, government policies and public health initiatives such as subsidy of electric bikes may facilitate greater uptake.

## 13 | CONCLUSION

E-cycling was associated with an increase in physiological response that can confer health benefits. The magnitude of effect in physiological responses was lower than compared with conventional cycling. Nonetheless, e-cycling is of sufficient intensity to meet PA recommendations. Compared with walking, e-cycling with electrical assistance offered greater physiological response. For many people, e-cycling offers an ideal entry approach for promotion of physical activity.

### ACKNOWLEDGEMENTS

We would like to acknowledge Ms Chrissy Freestone (Library Liaison, Deakin University) for their contribution to our study.

### CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

### AUTHOR CONTRIBUTIONS

JM and MK independently screened and reviewed titles and abstracts to identify eligible studies. JM, AB, and LP assessed full-text articles for eligibility. JM and RD independently performed data extraction and quality assessment for the studies included within the meta-analysis. JM conducted meta-analysis. Any discrepancies faced throughout the development of the manuscript were discussed with RM. All authors contributed to the writing of manuscript. All authors read and approved final manuscript.

### DATA AVAILABILITY STATEMENT

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

### ORCID

Jenna McVicar  <https://orcid.org/0000-0001-5007-7223>

### REFERENCES

1. Caprara G. Mediterranean-type dietary pattern and physical activity: the winning combination to counteract the rising burden of non-communicable diseases (NCDs). *Nutrients*. 2021;13(2):429.
2. Bull FC, Al-Ansari SS, Biddle S, et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med*. 2020;54(24):1451-1462.
3. Ding D, Lawson KD, Kolbe-Alexander TL, et al. The economic burden of physical inactivity: a global analysis of major non-communicable diseases. *Lancet*. 2016;388(10051):1311-1324.
4. Hoare E, Stavreski B, Jennings GL, Kingwell BA. Exploring motivation and barriers to physical activity among active and inactive Australian adults. *Sports*. 2017;5(3):47.

5. Shaw C, Keall M, Guiney H. What modes of transport are associated with higher levels of physical activity? Cross-sectional study of New Zealand adults. *Journal of Transport & Health*. 2017;7:125-133. doi:10.1016/j.jth.2017.09.010
6. Van Cauwenberg J, de Geus B, Deforche B. Cycling for transport among older adults: Health benefits, prevalence, determinants, injuries and the potential of E-bikes. In: Curl A, Musselwhite C, eds. *Geographies of Transport and Ageing*. Springer; 2018:133-151. doi:10.1007/978-3-319-76360-6\_6
7. Sundfør HB, Fyhri A. A push for public health: the effect of e-bikes on physical activity levels. *BMC Public Health*. 2017;17(1):1-12.
8. Bourne JE, Cooper AR, Kelly P, et al. The impact of e-cycling on travel behaviour: a scoping review. *J Transp Health*. 2020;19:100910.
9. Bourne JE, Sauchelli S, Perry R, et al. Health benefits of electrically-assisted cycling: a systematic review. *Int J Behav Nutr Phys Act*. 2018;15(1):1-15.
10. Peterman JE, Morris KL, Kram R, Byrnes WC. Pedelects as a physically active transportation mode. *Eur J Appl Physiol*. 2016;116(8):1565-1573.
11. Project EPHPP. *Quality Assessment Tool for Quantitative Studies*. McMaster University, School of Nursing. <https://merst.ca/ephpp/>
12. Ubago-Guisado E, Gracia-Marco L, Cavero-Redondo I, et al. Effect of different types of exercise on health-related quality of life during and after cancer treatment: a protocol for a systematic review and network meta-analysis. *BMJ Open*. 2019;9(11):e031374.
13. Sawilowsky SS. New Effect Size Rules of Thumb. *Journal of Modern Applied Statistical Methods*. 2009;8(2):597-599. doi:10.22237/jmasm/1257035100
14. Bini R, Wundersitz D, Kingsley M. Biomechanical and physiological responses to electrically assisted cycling during simulated mail delivery. *Appl Ergon*. 2019;75:243-249.
15. Hansen D, Soors A, Deluyker V, Frederix I, Dendele P. Electrical support during outdoor cycling in patients with coronary artery disease: impact on exercise intensity, volume and perception of effort. *Acta Cardiol*. 2018;73(4):343-350.
16. Langford BC, Cherry CR, Bassett DR, Fitzhugh EC, Dhakal N. Comparing physical activity of pedal-assist electric bikes with walking and conventional bicycles. *Journal of Transport & Health*. 2017;6:463-473. doi:10.1016/j.jth.2017.06.002
17. Louis J, Brisswalter J, Morio C, Barla C, Temprado J-J. The electrically assisted bicycle: an alternative way to promote physical activity. *Am J Phys Med Rehabil*. 2012;91(11):931-940.
18. Simons M, Vanes E, Hendriksen I. Electrically Assisted Cycling. *Medicine & Science in Sports & Exercise*. 2009;41(11):2097-2102. doi:10.1249/mss.0b013e3181a6aaa4
19. Alessio HM, Reiman T, Kemper B, von Carlowitz W, Bailer AJ, Timmerman KL. Metabolic and Cardiovascular Responses to a Simulated Commute on an E-Bike. *Translational Journal of the American College of Sports Medicine*. 2021;6(2):e000155. doi:10.1249/tjx.0000000000000155
20. Sperlich B, Zinner C, Hébert-Losier K, Born D-P, Holmberg H-C. Biomechanical, cardiorespiratory, metabolic and perceived responses to electrically assisted cycling. *Eur J Appl Physiol*. 2012;112(12):4015-4025.
21. La Salle DT, Shute R, Heesch M, Slivka D. Demands of simulated commuting using an electrically assisted bicycle. *Int J Exerc Sci*. 2017;10(3):454.
22. Hoj TH, Bramwell JJ, Lister C, et al. Increasing active transportation through e-bike use: pilot study comparing the health benefits, attitudes, and beliefs surrounding e-bikes and conventional bikes. *JMIR Public Health Surveill*. 2018;4(4):e10461.
23. Hall C, Hoj TH, Julian C, et al. Pedal-assist mountain bikes: a pilot study comparison of the exercise response, perceptions, and beliefs of experienced mountain bikers. *JMIR Form Res*. 2019;3(3):e13643.
24. Gojanovic B, Welker J, Iglesias K, Daucourt C, Gremion G. Electric bicycles as a new active transportation modality to promote health. *Med Sci Sports Exerc*. 2011;43(11):2204-2210.
25. Theurel J, Theurel A, Lepers R. Physiological and cognitive responses when riding an electrically assisted bicycle versus a classical bicycle. *Ergonomics*. 2012;55(7):773-781.
26. Meyer D, Steffan M, Senner V. Impact of Electrical Assistance on Physiological Parameters During Cycling. *Procedia Engineering*. 2014;72:150-155. doi:10.1016/j.proeng.2014.06.026
27. Berntsen S, Malnes L, Langåker A, Bere E. Physical activity when riding an electric assisted bicycle. *Int J Behav Nutr Phys Act*. 2017;14(1):1-7.
28. Fyhri A, Fearnley N. Effects of e-bikes on bicycle use and mode share. *Transportation Research Part D: Transport and Environment*. 2015;36:45-52. doi:10.1016/j.trd.2015.02.005
29. Fishman E, Cherry C. E-bikes in the mainstream: reviewing a decade of research. *Transp Rev*. 2016;36(1):72-91.
30. de Kruijf J, Ettema D, Dijst M. A longitudinal evaluation of satisfaction with e-cycling in daily commuting in the Netherlands. *Travel Behaviour and Society*. 2019;16:192-200. doi:10.1016/j.tbs.2018.04.003
31. Hausteijn S, Møller M. Age and attitude: changes in cycling patterns of different e-bike user segments. *Int J Sustain Transp*. 2016;10(9):836-846.
32. Kazemzadeh K, Koglin T. Electric bike (non) users' health and comfort concerns pre and peri a world pandemic (COVID-19): A qualitative study. *J Transp Health*. 2021;20:101014.
33. The UK Electric Bike Company Ltd. eBike To Work Scheme. [www.e-bikeshop.co.uk](http://www.e-bikeshop.co.uk)
34. Bieliński T, Ważna A. Electric scooter sharing and bike sharing user behaviour and characteristics. *Sustainability*. 2020;12(22):9640.
35. Murtagh EM, Nichols L, Mohammed MA, Holder R, Nevill AM, Murphy MH. The effect of walking on risk factors for cardiovascular disease: an updated systematic review and meta-analysis of randomised control trials. *Prev Med*. 2015;72:34-43.
36. van Namen M, Prendergast L, Peiris C. Supervised lifestyle intervention for people with metabolic syndrome improves outcomes and reduces individual risk factors of metabolic syndrome: a systematic review and meta-analysis. *Metabolism*. 2019;101:153988.
37. Wahid A, Manek N, Nichols M, et al. Quantifying the association between physical activity and cardiovascular disease and diabetes: a systematic review and meta-analysis. *J Am Heart Assoc*. 2016;5(9):e002495.
38. Manaugh K, Boisjoly G, El-Geneidy A. Overcoming barriers to cycling: understanding frequency of cycling in a University setting and the factors preventing commuters from cycling on a regular basis. *Transportation*. 2017;44(4):871-884.

39. Fyhri A, Heinen E, Fearnley N, Sundfør HB. A push to cycling—exploring the e-bike's role in overcoming barriers to bicycle use with a survey and an intervention study. *Int J Sustain Trans.* 2017;11(9):681-695.
40. McQueen M, MacArthur J, Cherry C. The E-Bike Potential: Estimating regional e-bike impacts on greenhouse gas emissions. *Transportation Research Part D: Transport and Environment.* 2020;87:102482. doi:10.1016/j.trd.2020.102482

**How to cite this article:** McVicar J, Keske MA, Daryabeygi-Khotbehsara R, Betik AC, Parker L, Maddison R. Systematic review and meta-analysis evaluating the effects electric bikes have on physiological parameters. *Scand J Med Sci Sports.* 2022;00:1–13. doi:[10.1111/sms.14155](https://doi.org/10.1111/sms.14155)

## SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.